

HemaClear® Placement

NO-HEMACLEAR® ZONE ARM

Ulnar Nerve, Radial Nerve,
Elbow, Forearm Muscles



NO-HEMACLEAR® ZONE LEG

Peroneal Nerve, Knee,
Calf Muscles -
Soleus/Gastrocnemius



OPTIMAL POSITION - UPPER ARM

Bottom of Deltoid Muscle
HemaClear®
Medium Yellow or Large Orange



OPTIMAL POSITION - FOREARM

10 cm Above the Wrist
Distal to Forearm Muscles
HemaClear® Model F™



OPTIMAL POSITION - THIGH

As Close to the Groin as Possible
HemaClear® Extra-Large Black & White
or Large Orange or Large Brown



OPTIMAL POSITION - LOWER LEG

10-15 cm Above the Lateral Malleolus
HemaClear® Model A™
or Large Orange



- Do not use **HemaClear®** on patients with poor peripheral blood flow, edema, or Deep Vein Thrombosis (DVT). See the **Wells Score System*** for likelihood of DVT.
- Do not use **HemaClear®** if the limb is infected or with malignancy.
- Do not apply **HemaClear®** directly on skin that is fragile or has significant lesions. Use a sterile ace bandage to protect fragile skin before applying **HemaClear®**.
- **Do not leave HemaClear® on a patient's limb for more than 120 minutes.**
- Do not place **HemaClear®** directly over the ulnar nerve (at the elbow) or the peroneal nerve (at the proximal tibia).

* *Wells PS, Anderson DR, Rodger M et al. (2003). Evaluation of D-dimer in the diagnosis of a suspected deep-vein thrombosis. New England Journal of Medicine 349: 1227–35.*